

Meeting Content
Message from the Division
Slide 1

Title

The Indiana Gypsy Moth Management Program depends on public participation.

Public participation in the Management program is necessary because ultimately either gypsy moth infestations or the actions taken to control them will affect the public. To be able to actively participate the Division holds several public meetings in areas where gypsy moth has been detected but is not yet causing damage. When gypsy moth is first detected the public may not be aware that there is a problem, that there will be a problem or what can or is being done to alleviate the problem.

The main purpose of the public meeting is to involve the public in the Gypsy moth management process by

Providing information

About gypsy moth biology

What damage can the gypsy moth cause?

How are gypsy moth infestations detected?

What is Slow the Spread Program?

What are the options for control?

What does the DNR propose?

There are several Governmental agencies cooperating in this effort.

Indiana Department of Natural Resources

- Division of Entomology and Plant Pathology – Public Outreach and Regulatory, and Control
- Division of Forestry – Survey, detection and control
- US Forest Service – Slow the Spread Project Coordination and Control
- USDA Animal and Plant Health Inspection Service – Plant Protection and Quarantine Eradication
- Purdue University – Public Outreach and Information, research

The public is encouraged to participate by commenting on how they want proceed with Gypsy moth management in their communities.

The division has many proposals and tools available to assist the communities but it is ultimately the decision of the public that will determine what if any action will be taken.

Participation does not end at this meeting. Comments will be received until March 14th when a final decision will need to be made for each location where action is being proposed.

Slide 2

Objectives

The objectives of the meeting are to provide information on the following

Biology

Damage

Survey/Maps

Treatment Options

DNR proposal

Questions and Comments

Slide 3

Life Stage Preview

Biology – The gypsy moth is an insect that goes through 4 life stage changes egg, larva, pupa, and adult.

Slide 4

Egg mass

The egg is laid in late summer and over winters. It hatches in early May in northern Indiana. There are anywhere from 500 – 1000 eggs in an egg mass. Eggs can be laid anywhere.

Slide 5

Early instar

After they hatch they move to the foliage. If it is crowded they spin a silk thread and balloon or blow to another tree or hilltop. Then they start to feed.

Slide 6

Ballooning

Slide 7

Late instar

As they feed they grow. Each growth spurt is called an instar. The male larvae have 5 instars the female have 6. Early instars feed during the day. Late instars feed at night. They crawl down from the foliage and find shelter during the day. The last instar feeds 24 hours. A late instar caterpillar can eat a square foot of leaves in a 24-hour period. They can eat 11 square feet of foliage during their entire caterpillar stage.

Slide 8

Caterpillars at the base of a tree

Slide 9

Pupae

After eating they change into a pupa. They attach themselves to a stable object to pupate. This can be on a tree trunk or the bumper of your car. They do not spin a cocoon. The pupae are bare except for small, tan hairs.

Slide 10

Adults

Male moths emerge from their pupal cases first in early July. The females emerge three to five days later. The males have feathery antennae, are dark brown with an inverted V mark pointing to a dot on their wings. Females have similar wing markings but are light in color, are larger, do not have feathery antennae and can't fly. Each female produces a pheromone that attracts the male to her. They mate and she lays eggs in masses.

Slide 11

She will lay eggs on most anything. The eggs over winter and the cycle starts again.

Slide 12

Damage: Why worry about gypsy moth?

Gypsy moth is not native to the United States

Only a few of its natural enemies have become established in the United States

Populations increase quickly. Each female lays 500 – 1000 eggs half of which may be females.

Caterpillars have a wide host range – oaks are their favorite

They are easily distributed naturally and by people

They can adversely affect the health of people and plants

They have an environmental impact that can be negative or positive

Slide 13

Hosts

Slide 14

Defoliation:

Reduces a tree's ability to produce and store food

The tree begins to decline

Branches and limbs begin to die

The tree becomes more susceptible to attacks from diseases and insects

Death may occur after just 2 years of 50% defoliation

Slide 15

Forests:

Decreased vigor and decline of host trees

Reduces Timber value

Defoliation opens the forest canopy and exposes the forest floor

Species composition change occurs when dead trees are replaced by hosts less favored by the gypsy moth

Forest floor dries in spring. Seed germination may decline

Dead limbs and trees increase the fire load – forest fires spread faster

Reduces recreational uses and value

Slide 16

Defoliation picture

Slide 17

Wildlife:

Hard mast will decrease (acorns, hickory nuts). It will eventually recover but at low levels in a forest where species composition has changed.

Soft mast will increase. Some wildlife species will utilize soft mast.

Bird populations fluctuate. An open canopy in spring predisposes nests to predation and parasitism (cow birds)

Food availability changes – some birds will utilize the gypsy moth caterpillars for food.

Others that depend on native caterpillar species for food may disperse to non-defoliated areas.

Gypsy moths eat the foliage that some native caterpillars utilize thereby reducing the food base for native defoliators. Native butterfly and moth caterpillars may decline.

Mammal populations will increase or decrease depending on food availability

Species may increase or decrease.

Slide 18

Soil:

- An open canopy exposes soil to sun and rain
- Soil temperatures increase
- The soil dries quicker
- Erodes more readily dependant on slope. Erosion removes valuable nutrients that would help trees survive future defoliations
- Reduces leaf litter

Slide 19

Water:

- Soil washed into streams and lakes increases turbidity
- Soil in water holds heat longer thereby increasing water temperatures
- Aquatic weeds may grow faster, die and produce organic matter
- Nutrient loads from caterpillar excrement, run off and aquatic weeds will reduce oxygen levels in some lakes and ponds.
- Aquatic species diversity will be reduced.
- Some lakes and reservoirs will have more water because of increased runoff.

Slide 20

- Urban Forest:** Where trees are already stressed
- Homeowners are liable for fallen limbs, tree removal costs and replacement costs
- Property values go down
- Ecological damage
- Trees cool the air in the summer, block wind, buffer noise, clean the air, and provide homes and shelter for wildlife. If removed these ecological benefits will be lost.

Slide 21

Larvae

- Thousands of crawling, smelling caterpillars.

Slide 22

Droppings

- Their smell and their fecal pellets fall from the trees.

Slide 23

Caterpillar hairs

- Drifting caterpillar hairs cause some Public health concern. There may be skin and respiratory reaction to the erticating hairs.

Slide 24

Survey – Why survey for Gypsy moth?

- Several states, along the leading edge of the infestation, trap for the male moths.
- Indiana sets traps on a grid system across the entire State

Slide 25

Both negative and positive trap catches give us valuable information:
accidental introductions
infestations
population movement
treatment evaluations

Slide 26

There are two types of traps:

Detection and monitoring

They use female sex pheromone to attract a male. The male then gets caught in the trap and is counted.

Monitoring trap holds 250 moths

Detection holds 30

Slide 27

By using trapping information we can divide the State into three zones

Non infested zone = detection

Transition zone = Slow the Spread action zone

Generally infested zone = monitoring zone

Generally infested – Suppression activities Gypsy moth is established and cannot be eradicated. Populations grow and coalesce. Defoliation is noticeable and damage begins to occur after two consecutive years of heavy defoliation. Natural movement (caterpillars ballooning) is frequent. Natural enemies may be present but are not at levels to maintain populations. This area is monitored to evaluate defoliation and outbreaks.

Transition zone – Slow the Spread Action Zone Populations of gypsy moth are small and isolated. They are either caused by caterpillars blowing from the generally infested area or life stages brought in by people. There is no noticeable defoliation. Damage is not evident but imminent. Natural enemies are usually not present but if present would not be in sufficient numbers to prevent damage. This area is trapped to locate populations, determine their size and growth potential. Some type of action is recommended in this area to delay establishment. No action will increase the size of the generally infested area significantly.

Non – infested – Eradication area. No male moths are caught or a few single moths are caught; no populations of gypsy moth have been detected. Natural movement of gypsy moth or effects of defoliation do not immediately threaten this area. Gypsy moths are introduced by human activity not by natural spread. There are no natural enemies in the area that would control gypsy moth. This area is trapped to detect infestations early. If an isolated infestation is found eradication is recommended.

Slide 28

Egg mass survey:

Based on trapping survey data we then select sites to look for egg masses

Trapping data, trapping history, egg mass finds or not, host availability, distance to the generally infested area, potential ecological threat all will help determine what we propose as an action.

Slide 29

Historical picture of egg mass survey and removal

Slide 30

To treat or not to treat?

Slide 31

Without slow the spread action

Slide 32

With slow the spread action

Slide 33

Slow the Spread

Indiana is in the program
Though gypsy moth will never be totally eradicated movement can be slowed down
New technology is becoming available
Slowing gypsy moth allows time for the natural enemies to naturally keep populations down over longer periods without human intervention.

Slide 34

Natural enemies are

Predators – eat gypsy moth life stages
Pathogens – are bacteria, fungus or virus that utilize gypsy moth for reproduction in the end killing gypsy moth life stages
Parasitoids – other insects that reproduce inside gypsy moth life stages eventually kill the host.

Slide 35

History of Natural enemies

They have been released over the years to help us fight populations of gypsy moth
More than 60 species have been introduced to the United States
To date only 11 parasitoids, 1 predator, and two pathogens have become established in North America
For the natural enemies to work gypsy moth populations can't move faster than the natural enemies so we try to control gypsy moth populations. Reduce long-term damage

Slide 36

What are the options?

DNR

Do nothing
Monitor
Mechanical mass trapping, burlap, egg mass treatment
Use biological pesticides Btk and Mating disruption (pheromone flakes)
Treatment options

Public

Do Nothing
Mechanical – burlap band
Treat egg masses with soybean oil
Treat trees with chemical insecticides

Treat with biological insecticides

The pheromone is not available for public use or application at this time.

Slide 37

What if we do nothing?

The county will be quarantined

Current counties: Allen, Dekalb, Noble, Steuben, Lagrange, Elkhart, Porter

Regulated articles are inspected to prevent movement of gypsy moth life stages.

Damage will occur sooner.

Slide 38

Proposed Action based on site evaluation

Slow the Spread using B.t.k., Pheromone flakes or both

Slide 39

Conditions considered when proposing a treatment

Mass trap when a few moths are caught in an area and there is no other trapping history.

Btk when we find high moth catch and other life stages are present

Mating disruption (pheromone flakes) high moth catch but no other life stages found

Hosts in the area

Distance from the generally infested area

Potential ecological threat

Slide 40

What is the material we are proposing to use and how is it applied

***Bacillus thuringiensis* variety *kurstaki* (Btk)** is a natural occurring soil bacteria.

When the caterpillar ingests it the Btk forms a crystal in the caterpillars gut. Not

all caterpillars are as susceptible as gypsy moth but it only affects caterpillars. It

is nontoxic to humans and other mammals and is most efficiently applied from the

air.

Slide 41

Mating disruption (pheromone flakes)

Female sex pheromone is distributed throughout the block before the males

emerge. Males emerge first but can't locate a specific female because he can't

track a single scent and he gives up. When the females emerge the males won't

mate. This only works in low level populations

Slide 42

Security

Material will be secured before during and after the operation

The plane will be secured

The DNR will have personnel monitoring from the ground and at the airport

Slide 43

Trapping will follow the aerial operation to evaluate the treatment and add additional population pressure.

Comments

Division of Entomology and Plant Pathology
402 W Washington Street, Room W 290
Indianapolis, IN 46204
317 – 232- 4120
317-232-2649
Email gjansen@dnr.state.in.us
www.state.in.us/dnr/entomolo

Questions About Gypsy moth

Comments